

# MATH 2554 : Cheat Sheet (No, you can't bring me to the exam)

## Nifty rules

### Derivation

1.  $\frac{d}{dx}c = 0$
2.  $\frac{d}{dx}f(x) + g(x) = f'(x) + g'(x)$
3.  $\frac{d}{dx}f(x)g(x) = f'(x)g(x) + f(x)g'(x)$
4.  $\frac{d}{dx}x^n = nx^{n-1}$
5.  $\frac{d}{dx}cf(x) = cf'(x)$
6.  $\frac{d}{dx}f(x) - g(x) = f'(x) - g'(x)$
7.  $\frac{d}{dx}\frac{f(x)}{g(x)} = \frac{g(x)f'(x) - f(x)g'(x)}{g(x)^2}$
8.  $\frac{d}{dx}f(g(x)) = f'(g(x)) \cdot g'(x)$

The above show the following rules : constant rule (1), constant multiple rule (5), sum rule (2 & 6), product rule (3), quotient rule (7), power rule (4), chain rule (8)

### Integration

1.  $\int \frac{1}{x} = \ln|x| + C$
2.  $\int f(g(x))g'(x)dx = \int f(u)du$
3.  $\int x^n dx = \frac{x^{n-1}}{n-1} + C$
4.  $\int_a^b f(g(x))g'(x)dx = \int_{g(a)}^{g(b)} f(u)du$

The above show the following rules : power rule (3), an exception to the power rule (1), substitution rule (2 & 4)

**Intermediate Value Theorem :** Suppose  $f$  is continuous on the interval  $[a, b]$  and  $L$  is a number strictly between  $f(a)$  and  $f(b)$ . Then there exists at least one number  $c$  in  $(a, b)$  satisfying  $f(c) = L$

### Definition of the Derivative :

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

**Mean Value Theorem :** If  $f$  is continuous on the closed interval  $[a, b]$ , then there is at least one point  $c$  in  $a, b$  such that

$$\frac{f(b) - f(a)}{b - a} = f'(c)$$

Note that **Rolle's Theorem** is a special case of MVT where  $f(a) = f(b)$

**Linear Approximation to  $f$  at  $a$  :** Suppose  $f$  is differentiable on an interval  $I$  containing the point  $a$ . The linear approximation to  $f$  at  $a$  is the linear function

$$L(x) = f(a) + f'(a)(x - a)$$

**L'Hopital's Rule :** Suppose  $f$  and  $g$  are differentiable on an open interval  $I$  containing  $a$  with  $g'(x) \neq 0$  when  $x \neq a$ . If  $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ , then

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

**Fundamental Theorem of Calculus :** If  $f$  is continuous on  $[a, b]$  and  $F$  is any antiderivative of  $f$  on  $[a, b]$ , then

$$(1) \quad F(x) = \int_a^x f(t)dt$$

$$(2) \quad \int_a^b f(x)dx = F(b) - F(a)$$

**Average Value of a Function :**

$$\bar{f} = \frac{1}{b-a} \int_a^b f(x) dx$$

### Basic derivative forms

**Trig derivatives :**

$$1. \frac{d}{dx} \sin x = \cos x$$

$$3. \frac{d}{dx} \tan x = \sec^2 x$$

$$5. \frac{d}{dx} \sec x = \sec x \tan x$$

$$2. \frac{d}{dx} \cos x = -\sin x$$

$$4. \frac{d}{dx} \cot x = -\csc^2 x$$

$$6. \frac{d}{dx} \csc x = -\csc x \cot x$$

**Inverse Trig derivatives :**

$$1. \frac{d}{dx} \arcsin x = \frac{1}{\sqrt{1-x^2}}$$

$$3. \frac{d}{dx} \arctan x = \frac{1}{1+x^2}$$

$$5. \frac{d}{dx} \operatorname{arcsec} x = \frac{1}{|x|\sqrt{x^2-1}}$$

$$2. \frac{d}{dx} \arccos x = -\frac{1}{\sqrt{1-x^2}}$$

$$4. \frac{d}{dx} \operatorname{arccot} x = -\frac{1}{1+x^2}$$

$$6. \frac{d}{dx} \operatorname{arccsc} x = -\frac{1}{|x|\sqrt{x^2-1}}$$

**Exponential/Log derivatives :**

$$1. \frac{d}{dx} e^x = e^x$$

$$3. \frac{d}{dx} b^x = b^x \ln b$$

$$2. \frac{d}{dx} \ln |x| = \frac{1}{x}$$

$$4. \frac{d}{dx} \log_b |x| = \frac{1}{x \ln b}$$

**...And for integrals, all these backwards!**

### Other

**Limits**

$$1. \lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

$$2. \lim_{x \rightarrow 0} \frac{\cos x - 1}{x} = 0$$

**Deriving Antiderivatives :**

$$1. \frac{d}{dx} \int_a^x f(t) dt = f(x)$$

$$3. \frac{d}{dx} \int_x^b f(t) dt = -f(x)$$

$$2. \frac{d}{dx} \int_a^{g(x)} f(t) dt = f(g(x))g'(x)$$

$$4. \frac{d}{dx} \int_{h(x)}^{g(x)} f(t) dt = f(g(x))g'(x) - f(h(x))h'(x)$$

**Common integrals utilizing substitution**

$$1. \int \cos ax dx = \frac{1}{a} \sin ax + C$$

$$7. \int \sin ax dx = -\frac{1}{a} \cos ax + C$$

$$2. \int \sec^2 ax dx = \frac{1}{a} \tan ax + C$$

$$8. \int \csc^2 ax dx = -\frac{1}{a} \cot ax + C$$

$$3. \int \sec ax \tan ax dx = \frac{1}{a} \sec ax + C$$

$$9. \int \csc ax \cot ax dx = -\frac{1}{a} \csc ax + C$$

$$4. \int e^{ax} dx = \frac{1}{a} e^{ax} + C$$

$$10. \int b^x dx = \frac{1}{\ln b} b^x + C$$

$$5. \int \frac{dx}{a^2 + x^2} = \frac{1}{a} \arctan \frac{x}{a} + C$$

$$11. \int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a} + C$$

$$6. \int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \operatorname{arcsec} \left| \frac{x}{a} \right| + C$$